

STUDIES OF ELECTROSTATIC WAVES, VLF-WAVE PARTICLE INTERACTIONS, AND PROPAGATIONS IN THE IONOSPHERE

Paul M. Kintner
School of Electrical Engineering
302 Rhodes Hall
Cornell University
Ithaca, NY 14853

phone: (607) 255-5304, fax: (607) 255-6236, e-mail: paul@ee.cornell.edu

Award #: N00014J921822

LONG-TERM GOAL

The long-term goal of this project is to study waves, irregularities, and propagation in the earth's ionosphere, particularly with respect to GPS. The project has two foci: first, to understand how the ionosphere affects GPS signals and second, how GPS can be used for remote sensing of the ionosphere. In association with these goals, we additionally conduct in situ experiments on satellites and sounding rockets to directly measure ionospheric waves and irregularities that affect signal propagation.

SCIENTIFIC OBJECTIVES

The scientific objectives of the project are to:

1. Develop GPS receivers for measuring scintillations and scintillation effects on GPS signals and receivers;
2. Continue study of the effects of equatorial scintillation storms on GPS through field campaigns, deployment of GPS scintillation receivers at collaborating institutions, and collaboration with Boston University studies employing airglow imaging and GPS TEC;
3. Extend GPS scintillation measurements to high latitudes by fielding the Cornell GPS scintillation receiver at Poker Flat, Alaska and Longyearbyen, Svalbard;
4. Develop new instrumentation for in situ studies of ionospheric irregularities based on the principles of very small payloads (nanosatellites), sounding rocket schedules and cost targets, and multiple (clustered) satellite missions; and
5. Develop an electric field experiment for an Indian satellite with the scientific goal of examining the possibility of ELF and VLF wave activity being associated with earthquakes.

Our research focuses on the study of waves, irregularities, and wave-particle interactions in space plasmas as well as the effects of these processes on radio signals. The vast majority of the universe exists in a plasma state, including our own upper atmosphere, or ionosphere, which is both a natural laboratory for studying space physics and a part of our environment affecting satellites and their signals. If the plasma is collisionless, which applies to our own ionosphere above roughly 150 km altitude, energy exchange and plasma acceleration must be produced by electric fields, usually in the form of plasma waves. The process of energy exchange and plasma acceleration mediated by plasma waves is termed wave-particle interactions.

For many years, wave-particle interactions implied the study of whistler mode waves interacting with radiation belt electrons on closed geomagnetic field lines. Our study is much more general

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE Studies of Electrostatic Waves, VLF-wave Particle Interactions, and Propagations in the Ionosphere			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Cornell University,School of Electrical Engineering,302 Rhodes Hall,Ithaca,NY,14853			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

and includes the transverse acceleration of ions, the interaction of ion beams with the ionosphere, the production of auroral electrons, the production of waves by auroral electron beams, and the production and transport of ionospheric irregularities, including those producing scintillations.

APPROACH

We are known for developing unique and ground-breaking experiments and instruments. Our most widely known contribution has been the plasma wave interferometer, first developed on sounding rockets and then later included on such spacecraft as Viking, Freja, Polar, Fast, and Cluster. Another example of new instrumentation developed at Cornell is the intelligent snapshot receiver, which is now popular among many groups and was part of the Freja plasma wave experiment. More recently we have developed GPS receivers, both for making scintillation measurements and for providing time transfer capability on spacecraft.

Our scientific strategy emphasizes experimental development. We have chosen this route because the field of space science, especially the electrical properties of space, is still experimentally limited. Theories of space physics and space plasma physics are quite plentiful, but discriminating measurements are few and far between. Within this context one may well ask what areas need the most attention. The answer concerns nonlinear problems involving plasma waves and electric fields in collisionless environments and turbulent media. Incidentally, these areas are also examples that, at one extreme, can test theories of basic plasma physics, and at the other extreme, are important for the development and application of new communication and navigation technologies.

WORK COMPLETED

Completed works include the following:

1. Developed a GPS receiver for measuring scintillations;
2. Conducted three field campaigns at the magnetic equator (Ancon, Peru) and one field campaign in Poker Flat, Alaska to measure GPS scintillations using the Cornell GPS scintillation receiver;
3. Demonstrated that GPS scintillations could be used for measuring ionospheric irregularity drifts at the equator;
4. Demonstrated that velocity resonance could occur between irregularity drifts and GPS signal ionospheric penetration points, thereby greatly lengthening fade periods with major implications for GPS receiver design;
5. Demonstrated that a new kind of wave-particle interaction called lower hybrid solitary structures (which we had previously discovered) are examples of plasma self-organization containing very large, rotational electric fields and producing transverse ion acceleration. Currently we (and many international scientists) are investigating these structures as the source of O^+ ions for the magnetospheric radiation belts;
6. Published 12 papers on ionospheric irregularities and plasma waves, which were supported by ONR;
7. Delivered 3 invited reviews and 13 contributed papers at AGU annual meetings and international workshops;

8. Awarded a Ph.D. to one graduate, who subsequently accepted a Director's Fellowship with Los Alamos National Laboratory; and
9. Received a Cornell Teaching award for developing a GPS design course in engineering.

RESULTS

We demonstrated that velocity matching between equatorial spread-F irregularity drifts and GPS signal ionospheric penetration points greatly lengthen scintillation fade periods from seconds to tens of seconds or longer, with major implications for GPS receiver design. Although the velocity matching condition was demonstrated for a stationary receiver, it is not only expected to occur for mobile receivers, but is more likely to occur. This particularly applies to aviation receivers.

ONR funding of Cornell satellite instruments, particularly for Viking and Freja, along with NASA funding of Cornell sounding rocket experiments, has recently resulted in a major breakthrough in understanding high-latitude ion acceleration, the mass source for the magnetosphere. Of particular interest is that this breakthrough was made possible by a collaboration with NRL theorist Ganguli and laboratory experimentalists Walker and Amattuci. The critical breakthrough was the realization that the homogeneous theories were not applicable to the space environment and non-homogeneous models must be employed.

IMPACT/APPLICATION

The demonstration of velocity matching between irregularity drifts and GPS signal ionospheric penetration points and the subsequent discovery of lengthened scintillation fade periods has major implications for GPS receiver design. Since GPS is a CDMA system, which depends on code correlation to maintain signal tracking, we expect that during periods of velocity matching there will be a greatly increased rate of signal tracking failure, loss of lock, and lengthened acquisition times. Although the velocity matching condition was demonstrated for a stationary receiver, it is not only expected to occur for mobile receivers, but is more likely to occur. This particularly applies to aviation receivers. Where GPS integrity is an issue, this result has major implications.

TRANSITIONS

We will collect GPS scintillation data from several sites across South America to create a regional study of equatorial scintillations. We have selected four sites, chosen primarily because they have existing universities or institutes with ionosphere experts who can operate Cornell GPS scintillation receivers. The sites are:

1. Geophysical Institute of Peru (IGP) at Ancon, Peru;
2. National University at Tucuman, Argentina;
3. National Institute of Space Investigations (INPE) at Sao Paulo, Brazil; and
4. Northern Federal University at Natal, Brazil.

At each site, a senior scientist has been recruited for collaboration who also attended a workshop on GPS scintillations at Cornell this past Spring. Cornell GPS scintillation receivers were provided to each institution, who will operate the receivers on a campaign basis.

Additionally we will develop autonomous GPS receivers to conduct GPS scintillation measurements on a continuous basis, as opposed to operating in a campaign mode.

RELATED PROJECTS

One should note that constructing a regional model of equatorial scintillation storms and spread F is a major undertaking and that the chain of Cornell GPS scintillation receivers can only be one aspect of this ambitious objective. We will closely coordinate our work with Boston University, which has similar scientific objectives. They will be collecting and analyzing GPS TEC data from the JPL chain receivers at Arequipa, Bogata, Fortaleza, and Santiago. In addition, BU will be operating an airglow imager at Arequipa, Tucuman.

REFERENCES

S.P. Powell and P.M. Kintner, 1996. "Using GPS for telemetry system clock generation and synchronization on-board sounding rockets," Proceedings of ION GPS-96, The 9th International Technical Meeting of *The Satellite Division of The Institute of Navigation*, published by The Institute of Navigation, Alexandria, Virginia, part 2, 1341-1348.

T.L. Beach, M.C. Kelley, P.M. Kintner, and C.A. Miller, 1997. "Total electron content variations due to nonclassical traveling ionospheric disturbances: Theory and Global Positioning System observations," *J. Geophys. Res.*, 102(A4), 7279-7292.